

BIOMIN® 17% COPPER

SYSTEMIC

Product Information

Biomin® Copper is a mineral chelated mainly by amino acids. **Biomin® Copper** was designed for soil and / or foliar application. Completely bioavailable and non-phytotoxic to plants when applied in accordance to label directions.

Designed to prevent and correct Copper deficiencies and to boost crops during critical or fast growing periods.

Biomin® Copper can be applied to many crops including vegetables, potatoes, carrots, all berries and strawberries, brassicas, all fruit trees, avocados, persimmons, citrus, all nut trees, vineyards, kiwifruit, tropical fruit, almonds, all melons, beans, ginger, ornamentals, cut flowers and pastures.

GUARANTEED ANALYSIS

Dry Weight Basis %/w/w

As true amino acid Chelates:

COPPER (Cu) 17%

Copper is present as an amino acid chelate.

KEEP OUT OF REACH OF CHILDREN

PREVENTS AND CORRECTS MINERAL DEFICIENCIES OF COPPER

NET CONTENTS

☐ 2 kg ☐ 5 kg ☐ 10 kg ☐ 20 kg ☐ 25 kg

Lot #

Directions and General Recommendations

SOIL APPLICATION

Biomin® Copper may be applied before or after planting through irrigation water. The application rate is 0.5 - 3kg / Ha early in Spring, during the season and post harvest depending on crop, soil conditions and application methods. Repeat once more 3 - 4 weeks later. Do NOT exceed 5kg / Ha per season.

FOLIAR APPLICATION

DECIDUOUS FRUIT TREES (APPLES, PEARS & STONE FRUIT):

Apply **Biomin® Copper** at 1 - 2kg / Ha 2 -3 weeks before dormancy to assist with leaf defoliation and supply a systemic form of Copper into the crop.

GRAPES (WINE & TABLE GRAPES):

Apply **Biomin® Copper** at 0.5 - 1kg / Ha in early Spring when new shoots reach 20cm in length or when deficiency symptoms appear. Repeat the application if necessary 2 - 3 weeks later. Do NOT apply **Biomin® Copper** after fruit set.

COMPATIBILITY

To further increase availability, add **Biomin® Copper** to the tank when the pH of the water is 6.5 – 7.5. **Never** apply when pH is less than 6.5.

ALWAYS perform a compatibility test before mixing any products with **Biomin® Copper**.

WARNING

DO NOT APPLY **BIOMIN® COPPER** AS A FOLIAR APPLICATION ON COPPER SENSITIVE CROPS. FOR ANY CROPS NOT MENTIONED ON THIS LABEL CONTACT SUPPLIER BEFORE USING **BIOMIN® COPPER**

LIMITED WARRANTY

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Imported and Distributed in Australia exclusively by: Zadco For Quality Gro PTY LTD. ABN 57080 759 084 | 9/9A Foundry Road, Seven Hills NSW 2147
Ph: 02 9838 9111 Fax: 02 9838 9110 | email: inquiries@zadco.com.au website: www.zadco.com.au Imported and Distributed in New Zealand exclusively by:
Roots Shoots and Fruits Ltd | PO Box 72, Waiheke Island, Auckland, New Zealand | Ph: +64(0) 9372 9155 Fax: +64(0) 9372 9156 | email: rsf@rd2.co.nz
website: www.rd2.co.nz Manufactured by: JH BIOTECH, INC. PO Box 3538, Ventura, CA 93006, USA