

**ORGANIC**

# BIOMIN<sup>®</sup> N

## 5 - 0 - 0



### Product Information

#### About BIOMIN<sup>®</sup> N

BIOMIN<sup>®</sup> N is an organic liquid plant nutrient product derived from hydrolyzed soy protein. It is designed for soil and/or foliar applications.

BIOMIN<sup>®</sup> N is recommended for use on all crops including: alfalfa, almond, apple, apricot, avocado, banana, tropical fruits, barley, beans, carrot, cherry, citrus, corn, cotton, grape, garlic, hops, lettuce, brassicas, onion, peach, pear, pecan, pepper, pistachio, plum, potato, prune, rice, row crops, sorghum, soybean, strawberry, sugarbeet, sunflower, tomato, vegetable, walnut, and wheat.

#### Guaranteed Analysis

Total Nitrogen (N) 5.0%  
5% Water Soluble Nitrogen

*Derived from soy protein hydrolysate*

**KEEP OUT OF REACH OF CHILDREN**

**SHAKE WELL BEFORE USE  
DO NOT APPLY AT TIME OF BLOOM**

#### Net Contents

3.785L  18.92L  1000L

**LOT:  
BEST BEFORE:**

### Directions and General Recommendations

Dilute BIOMIN<sup>®</sup> N with at least 200 parts water and apply in accordance with the directions.

**VEGETABLES, LETTUCE, BRASSICAS, SUGARBEET, ROW CROPS, AND ROOT CROPS:** Apply 10 - 40L / ha at 2 week intervals when plants are well established depending on crop requirements.

**COMMERCIAL GREENHOUSES AND NURSERIES:** Inject into irrigation water at ratio of at least 1:1,000L. Apply at 2 week intervals.

**ALFALFA:** Apply 10 - 40L / ha one week after each cutting.

**APRICOTS, CHERRIES, HOPS, PLUMBS, PEACHES, GRAPES, APPLES, PECANS, PISTACHIOS, ALMONDS, PRUNES, WALNUTS, AND OTHER FRUITS AND NUTS:** Apply 10 - 40L / ha prior to bloom. Older trees may require another application 4-6 weeks later. Repeat the application 2 more times in 3-4 week intervals.

**BEANS, PEAS, SOYBEANS, TOMATOES, PEPPERS:** Apply 10 - 40L / ha at bud stage, prior to bloom. Repeat the application 2 more times in 3-4 week intervals.

**CITRUS, AVOCADOS, BANANA & TROPICAL FRUIT:** Apply 10 - 40L / ha at bud stage, prior to bloom. When new growth appears, repeat the application 2 times at 2-4 week intervals.

**COTTON:** Apply 10 - 40L / ha after true leaves appear. Second application should be made at early square at Apply 10 - 40L / ha. Repeat the application 2 more times at 3-4 week intervals.

**RICE, BARLEY, SORGHUM, WHEAT, AND OTHER SMALL GRAINS:** Apply 10 - 40L / ha at boot stage. Repeat the application prior to and after bloom.

**STRAWBERRY AND OTHER BERRIES:** Apply 10 - 40L / ha after transplanting. Repeat the application throughout the season at 2-4 week intervals.

**POTATOES, CARROTS, ONIONS & GARLIC:** Apply 10 - 40L / ha at tuber initiation. Repeat the application at 2-4 week intervals.

**CORN:** Apply 10 - 40L / ha prior to tasseling. Repeat the application at 2-4 week intervals.

**SUNFLOWER, CANOLA AND OTHER OILSEED CROPS:** Apply 10 - 40L / ha when plants are well established. Repeat the application at 2-4 week intervals.

#### Storage

Store BIOMIN<sup>®</sup> N in cool dry places. Avoid direct sunlight and heat. Dispose product / container in accordance with federal, state, and local regulations.

**PRECAUTIONARY STATEMENT:** do not swallow. Avoid contact with eyes.  
**FIRST AID:** *If swallowed:* seek medical attention. Rinse mouth. *If in eyes:* remove contact lenses if present. Wash thoroughly with water for several minutes.

#### Limited Warranty

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

**JHBiotech**  
A U S T R A L A S I A

Imported & Distributed in Australasia & NZ exclusively by: © JH Biotech Australasia Pty. Ltd.  
PO Box 6973 Baulkham Hills BC, Sydney NSW Australia 2153 | T (02) 8814 6245 | F (02) 8814 6247 | E info@jhbiotech.com.au  
Manufactured by JH BIOTECH, INC. PO Box 3538, Ventura, CA 93006, USA