

BIOMIN[®] TRACE BOOSTER

Zn 6% | Mg 4.1% | Fe 2.6% | Mn 2.6% | B 1.5% | Mo 0.5%

Product Information

About Biomin[®] Trace Booster

BIOMIN[®] TRACE BOOSTER is an amino acid chelated multi-mineral. **BIOMIN[®] TRACE BOOSTER** was designed for soil and/or foliar applications. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct trace mineral deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilisers. **BIOMIN[®] TRACE BOOSTER** can be applied to ALL crops including vegetables, potatoes, carrots, onions, all berries and strawberries, all brassicas, all fruit trees, mangos, avocados, persimmons, citrus, all nut trees, vineyards, kiwifruit, tropical fruit, lettuce, asparagus, all melons, beans, ginger, ornamentals, cut flowers, cereals and pastures.

GUARANTEED ANALYSIS

As true amino acid Chelates:	Dry Weight Basis %/w/w
Zinc (Zn)	6.0%
Magnesium (Mg)	4.1%
Iron (Fe)	2.6%
Manganese (Mn)	2.6%
Boron (B)	1.5%
Molybdenum (Mo)	0.5%
As amino acid:	
Nitrogen (N)	6.7%

Magnesium, Zinc, Iron & Manganese present as amino acid chelates. Molybdenum & Boron present as amino acid complexes. Nitrogen present as amino acid.

KEEP OUT OF REACH OF CHILDREN

PREVENTS AND CORRECTS MINERAL DEFICIENCIES

In conditions where the water and/or soil pH is alkaline, it would be advantageous to apply **MOBILIZER** to the water first before adding **BIOMIN[®] TRACE BOOSTER** to the tank. To further increase availability, add **BIOMIN[®] TRACE BOOSTER** to the tank when the pH of the water is 6.5 - 7.0.

Net Contents

2 Kg 10 Kg 25 Kg LOT #

Directions and General Recommendations

Soil Application

BIOMIN[®] TRACE BOOSTER may be applied before or at planting through irrigation water. The application rate is 2 - 3 kilograms per hectare depending on crop, soil conditions and application methods.

Foliar Application

ALL FIELD CROPS (grain crops, flowers, etc.): Apply 1.0 - 1.5 kilograms of **BIOMIN[®] TRACE BOOSTER** per hectare when crops are 15 - 30cms high or when deficiency symptoms appear. Repeat application 3 - 4 weeks later or as needed.

ALL VEGETABLE CROPS: Apply 1.0 - 1.5 kilograms of **BIOMIN[®] TRACE BOOSTER** per hectare at about 2 - 4 weeks after emergence or if deficiency symptoms appear. Repeat the application every 3 - 4 weeks, or as needed.

GREENHOUSE CROPS: Apply 100 - 150 grams of **BIOMIN[®] TRACE BOOSTER** per 100 liters of water 2-3 weeks after emergence or transplanting, or when trace element deficiencies first appear. Repeat the application every month.

APPLES: Apply 1.0 - 2.0 kilograms of **BIOMIN[®] TRACE BOOSTER** per hectare in early spring when new leaves emerge or 2 weeks before flowering. Repeat the application post-harvest at 6 weeks before dormancy at a rate of 1.0 - 2.0 kilograms per hectare.

STONE FRUIT, CHERRIES & ALMONDS: Apply 1.0 - 2 kilograms of **BIOMIN[®] TRACE BOOSTER**

Limited Warranty

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

per hectare at early pink bud formation. Repeat 3 - 4 weeks after fruit set. Repeat post-harvest at 2 kilograms per hectare.

ALL FRUIT TREES: Apply 1.0 - 2 kilograms of **BIOMIN[®] TRACE BOOSTER** per hectare in early spring when new leaves emerge or 2 weeks before flowering. Repeat mid-season. Repeat the application post-harvest at 6 weeks before dormancy.

KIWIFRUIT: Apply 1.0 - 2 kilograms of **BIOMIN[®] TRACE BOOSTER** per hectare in early spring when new leaves emerge or 2 weeks before flowering. Repeat the application post-harvest at 6 weeks before dormancy at a rate of 1.0 - 2.0 kilograms per hectare.

GRAPES, AVOCADOS & CITRUS: Apply 1.0 - 2 kilograms of **BIOMIN[®] TRACE BOOSTER** per hectare in early spring when new leaves emerge or 2 weeks before flowering or at any stage in the season when deficiency symptoms appear. Repeat the application post-harvest at a rate of 1.0 - 2.0 kilograms per hectare.

BIOMIN[®] TRACE BOOSTER is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 100 times of water.

Compatibility

BIOMIN[®] TRACE BOOSTER is NOT compatible with any Copper or Copper-containing products. Do NOT mix **BIOMIN[®] TRACE BOOSTER** with Phosphorus, Calcium or Potassium containing products. **BIOMIN[®] TRACE BOOSTER** is compatible with most pesticides. **ALWAYS run a jar test before application.**

ZADCO
for quality gro. pty. ld.

Manufactured by: JH BIOTECH, INC. PO
Box 3538, Ventura, CA 93006, USA

Imported and Distributed in Australia by: Zadco For Quality Gro PTY LTD. ABN 57080 759 084
PO BOX 6973 Baulkham Hills BC NSW 2153 Australia T: 02 8814 6638 | F: 02 8814 6623
E: inquiries@zadco.com.au | W: www.zadco.com.au

Imported & Distributed in New Zealand by: Roots Shoots & Fruits Ltd. | P.O. Box 72 Waiheke Island,
Auckland New Zealand T: 09-372-9155 | M: 027-6075-764 | F: 09-372-9156 | E: rsf@rd2.co.nz