# **FULZYME® PLUS**



# **Product Information**

## GENERAL INFORMATION

FULZYME PLUS is a natural product produced by a unique fermentation process. It contains a group of beneficial microorganisms and selected enzyme complexes specifically designed to improve soil and plant health by crowding out pathogens.

FULZYME PLUS can be applied to ALL crops including all vegetables, Potatoes, Carrots, Ohions, all berries and strawberries, all Brassicas, all Truit trees, Avocadoes, Persimonos, Cifrus, all Nut trees, Vineyards, Kiwlfruit, Tropical fruit, Lettuce, Asparagus, all Melons, Beans, Cilique, Omamentals, Cut flowers and Pastures.

#### GUARANTEED ANALYSIS

B. subtilis Bacteria 2 x 10 <sup>10</sup> / gram
 Amino Acids

Inert ingredients: Water

## KEEP OUT OF REACH OF CHILDREN

For increased benefits, combine FULZYME PLUS 1L/ha with TRI-025 (1kg/ha) and with HUMAX (1kg/ha).

COMPATIBILITY: FULZYME PLUS is compatible with TRI-D25 PHOSCARE and HUMAX.

FULZYME PLUS is INCOMPATIBLE with Copper, Chlorine, and or any fumigation products and anti bacterial products.

#### How to Clean Spray Tank, Filters, and Accessories:

Remove spray tank filters. Soak them for 3-4 hours in a concentrated trea solution (2kg of Urea in 10 It of water). Dissolve 5 kg of Urea in 1000Lt spray tank. Run the pump until all hoses and nozzles are filled with urea solution. Flush filters and spray tank prior to spraying FULZYME PLUS in the late afemoon.

#### NET CONTENTS

1 Litre

Lot # 2403311 Best before: April 2016

## Directions and General Recommendations

## SOIL APPLICATION

Apply FULZYME PLUS at 1L/ha in drip irrigation and at 2L/ha in Sprinkler or Pivot irrigation. Repeat six weeks later.

TRANSPLANTING DIP: Dilute FULZYME PLUS at 2 - 3mL / 1L of water and dip seedlings (roots and leaves) or water the seedlings while in trays before transplanting

POTTING MIXES: Dilute FULZYME PLUS at 50 - 100mL per cubic meter of potting mix.

ALL FIELD CROPS (grain crops, flowers, etc...): Apply 1.0 L per hectare of FULZYME PLUS at planting. Repeat the application four to six weeks later.

ALL VEGETABLE CROPS: Apply 1.0 L
per hectare of FULZYME PLUS at planting.
Repeat the application four to six weeks
later.

GREENHOUSE CROPS: Inject FULZYME PLUS at a rate of 100mL per shed (400 sqm). Repeat this application every month.

APPLES, POME & STONE FRUIT: Apply 1L per hectare of FULZYME PLUS at any stage in spring and Repeat early autumn. FRUIT TREES, NUTS & AVOCADOS: Apply 1L per hectare of FULZYME PLUS at any stage in the spring. Repeat mid season. Repeat early autumn.

KIWIFRUIT: Apply 1L per hectare of FULZYME PLUS at any stage in the spring. Repeat mid season. Repeat early autumn.

GRAPES & CITRUS: Apply 1L per hectare of FULZYME PLUS at any stage in the spring (OCT-NOV). Repeat early in autumn (March-Apr)

## FOLIAR APPLICATION

Activate **FULZYME PLUS** in water for 3 - 4 hours prior to spraying.

Apply FULZYME PLUS at 2 - 3 mL / 1L of water and spray on to the foliage and fruit of the plants early morning or late afternoon. Target applications towards the fruit on tree crops and bunches on grapes. Repeat as needed. No withholding period exists.

FULZYME PLUS is best applied when plants are in an active growing stage, after irrigation or rainfall. FULZYME PLUS may be used at any time throughout the season. For best results and more specific recommendations, contact your nearest distributor.

## LIMITED WARRANTY

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Imported and Distributed in Australia exclusively by: Zadoo For Ouality Core PTV.LTD. ABM 57 080 759 084 1 PO BDX.6935 BallAhrin Hills BC NSW 2158 7 Core 284 H6381 Face 284 H64281 Face 284 H