

HUMAX®



Product Information

HUMAX® is a humic acid product extracted from Leonardite and **HUMAX®** has been labelled the *Black Gold in Agriculture*. It consists of a mixture of macromolecules that may increase the uptake of mineral nutrients in plants. Humic acid is the major constituent of humus. **HUMAX®** promotes the conversion of a number of essential plant nutrients into available forms by chelating with the nutrients. **HUMAX®** and its derivatives, in small amounts, increase the permeability of cell membranes and hence promote the uptake of nutrients by plants. The colloidal nature of **HUMAX®** allows it to absorb a large quantity of water molecules and hence increases the water holding capacity of soil. **HUMAX®** can increase soil particle aggregation. As a result, it can increase soil aeration, tilth and workability, as well as better water movement.

HUMAX® can be applied to ALL crops including all vegetables, Potatoes, Carrots, Onions, all berries and strawberries, all Brassicas, all fruit trees, Avocadoes, Persimmons, Citrus, all Nut trees, Vineyards, Kiwifruit, Tropical fruit, Lettuce, Asparagus, all Melons, Beans, Ginger, Ornamentals, Cut flowers and Pastures.

GUARANTEED ANALYSIS

Humic Acid 80%
Derived from Leonardite

KEEP OUT OF REACH OF CHILDREN

HUMAX® is best dissolved by sprinkling the product over the top of warm water gently and allowing the product to be absorbed gradually by the water. It is not advisable to pour the entire contents of the product into water as it may make dissolving the product take longer than necessary.

COMPATIBILITY: Always perform a compatibility test before mixing any products with **HUMAX®**.

DO NOT mix **HUMAX®** with Calcium based products.

NET CONTENTS

5 kilograms 25 kilograms

Lot #

Directions and General Recommendations

SOIL APPLICATION

HUMAX® may be applied before or at planting through irrigation water. The application rate is 1.0 – 2.0 kilograms per hectare depending on crop, soil conditions and application methods. It can be applied to all crops as a soil application.

ALL FIELD CROPS (grain crops, flowers, pastures, etc...): Apply 1.0 – 2.0 kilograms of **HUMAX®** per hectare when crops are 15 - 30cms high or when deficiency symptoms appear. Repeat application as needed.

ALL VEGETABLE CROPS: Apply 1.0 – 2.0 kilograms of **HUMAX®** per hectare at about 2 - 4 weeks after emergence. Repeat the application as needed.

APPLES, POME & STONE FRUIT: Apply 1.0 – 2.0 kilograms of **HUMAX®** per hectare 3 - 4 times throughout the season. Reapply 6 weeks before dormancy also.

FRUIT TREES & AVOCADOS: Apply 1.0 – 2.0 kilograms of **HUMAX®** per hectare in early spring when new leaves emerge or 2 weeks before flowering. Repeat the application at least once every month throughout the season and also as a post harvest application.

KIWIFRUIT: Apply 1.0 – 2.0 kilograms of **HUMAX®** per hectare in early spring when new leaves emerge or 2 weeks before

flowering. Repeat the application at least once every month throughout the season and also as a post harvest application.

GRAPES: Apply 1.0 – 2.0 kilograms of **HUMAX®** per hectare in early spring when new leaves emerge or 2 weeks before flowering or any time. Repeat the application once every month throughout the season and also as a postharvest application.

FOLIAR APPLICATION

HUMAX® can be applied as a foliar application at a rate of 0.5 – 1.0 kilograms per hectare. **HUMAX®** should not be applied as a foliar application targeting fruit or flower crops after fruit set, especially on kiwifruit, apples, peaches, persimmons, nectarines, table grapes, strawberries or flowers. **MOBILIZER®** is a better alternative after fruit set for most crops as a foliar application.

HUMAX® is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 1000 times of water.

HUMAX® may be used at any time throughout the season. For best results and more specific recommendations, contact your nearest distributor.

LIMITED WARRANTY

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Imported and Distributed in Australia exclusively by: Zadco For Quality Gro PTY LTD. ABN 57080 759 084 | 9/9A Foundry Road, Seven Hills NSW 2147
Ph: 02 9838 9111 Fax: 02 9838 9110 | email: inquiries@zadco.com.au. website: www.zadco.com.au Imported and Distributed in New Zealand exclusively by:
Roots Shoots and Fruits Ltd | PO Box 72, Waiheke Island, Auckland, New Zealand | Ph: +64(0) 9372 9155 Fax: +64(0) 9372 9156 | email: rsf@rd2.co.nz
website: www.rd2.co.nz Manufactured by: JH BIOTECH, INC. PO Box 3538, Ventura, CA 93006, USA